

FINAL MATH 265

Evaluation out of 100. Calculators are NOT allowed.

1. (8) Find the volume of the solid enclosed by the parabolic cylinders $y = 1 - x^2$, $y = x^2 - 1$ and the planes $x + y + z = 2$, $2x + 2y - z + 10 = 0$.
2. (8) Use a double integral to find the area of the region within both of the circles $r = \sin \theta$ and $r = \cos \theta$ ((r, θ) are polar coordinates).
3. (8) The joint density function for a pair of random variables X and Y is

$$f(x, y) = Cx(1 + y) \text{ if } 0 \leq x \leq 1, \quad 0 \leq y \leq 2$$

and $f(x, y) = 0$ otherwise. Find a) the value of constant C ; b) $P(X \leq 1, Y \leq 1)$; c) $P(X + Y \leq 1)$

4. (8) Find the volume enclosed by the torus $\rho = \sin \varphi$ ((ρ, θ, φ) are spherical coordinates).
5. (8) Find the volume and centroid of the solid that lies above the cone $z = \sqrt{x^2 + y^2}$ and below the sphere $x^2 + y^2 + z^2 = 1$.
6. (8) Evaluate

$$\int_{-2}^2 \int_0^{\sqrt{4-y^2}} \int_{-\sqrt{4-x^2-y^2}}^{\sqrt{4-x^2-y^2}} y^2 \sqrt{x^2 + y^2 + z^2} dz dx dy$$

7. (8) Evaluate $\iint_R \cos \left\{ \frac{y-x}{y+x} \right\} dA$ using an appropriate change of variables, where R is the trapezoidal region with vertices $(1, 0)$, $(2, 0)$, $(0, 2)$ and $(0, 1)$.
8. (8) Find the work done by the force field $\mathbf{F}(x, y, z) = (y + z)\mathbf{i} + (x + z)\mathbf{j} + (x + y)\mathbf{k}$ on a particle that moves along the line segment from $(1, 0, 0)$ to $(3, 4, 2)$.
9. (8) Find the area of the helicoid $\mathbf{r}(u, v) = u \cos v \mathbf{i} + u \sin v \mathbf{j} + v \mathbf{k}$, $0 \leq u \leq 1$, $0 \leq v \leq \pi$
10. (8) Evaluate the surface integral $\iint_S \mathbf{F} d\mathbf{S}$ if $F = x\mathbf{i} + y\mathbf{j} + 2z\mathbf{k}$ and S is the surface of the solid bounded by hemisphere $x^2 + y^2 + z^2 \leq a^2$, $z \geq 0$ and the plane $z = 0$ with outward orientation.
11. (8) Evaluate $\iint_S \text{curl } \mathbf{F} d\mathbf{S}$ where

$$\mathbf{F}(x, y, z) = x^2 y z \mathbf{i} + y z^2 \mathbf{j} + z^3 e^{xy} \mathbf{k}$$

and S is the part of the sphere $x^2 + y^2 + z^2 = 5$ that lies above the plane $z = 1$ and S is oriented upwards.

12. (8) Solve the initial value problem

$$y'' + y' - 2y = x + \sin 2x, \quad y(0) = 1, \quad y'(0) = 0.$$

13. (6) A force of $400N$ stretches a spring 2 meters. A mass of 50 kilograms is attached to the end of the spring and released from the equilibrium position with an upward velocity of $10m/s$. Find position of the mass as function of time.